

*A multiplex set for microsatellite typing
and sexing of the European bee-eater
(Merops apiaster)*

**Ghaniya Bi, Natalie dos Remedios,
Deborah A. Dawson, Sjouke A. Kingma,
Julia Schroeder & Hannah L. Dugdale**

**European Journal of Wildlife
Research**

ISSN 1612-4642

Eur J Wildl Res
DOI 10.1007/s10344-016-1012-2



Your article is published under the Creative Commons Attribution license which allows users to read, copy, distribute and make derivative works, as long as the author of the original work is cited. You may self-archive this article on your own website, an institutional repository or funder's repository and make it publicly available immediately.

A multiplex set for microsatellite typing and sexing of the European bee-eater (*Merops apiaster*)

Ghaniya Bi¹ · Natalie dos Remedios¹ · Deborah A. Dawson¹ · Sjouke A. Kingma² · Julia Schroeder^{3,4} · Hannah L. Dugdale^{1,2,5}

Received: 16 November 2015 / Revised: 5 April 2016 / Accepted: 10 April 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract Microsatellite loci are widely used in ecological and evolutionary studies to assess inbreeding, genetic parentage and population structure. Such loci are often optimised in multiplexes to allow for economical and efficient use. Here, we tested 11 microsatellite loci designed for use in European bee-eaters (*Merops apiaster*), along with 31 loci isolated in other species, for their utility in European bee-eaters sampled on Susak Island, Croatia. Of these 42 loci, 20 were polymorphic in 38 individuals. These polymorphic loci were further assessed in a sub-set of 23 adults, excluding close relatives, and exhibited between three and 13 alleles each. All loci were autosomal, as indicated by the presence of heterozygotes in both males and females. One of the polymorphic loci exhibited low heterozygosity, three loci deviated from Hardy-Weinberg equilibrium and three pairs of loci displayed linkage disequilibrium. The remaining selected eight cross-species loci and seven loci isolated in European bee-eaters were combined with two sex-typing markers and optimised in five multiplexes. A combination of 15 autosomal loci of varying degrees of polymorphism makes this multiplex set particularly suitable for both parentage and spatial genetic analyses. This

multiplex set therefore provides a useful toolkit for studying kin selection and population genetics in the cooperatively breeding European bee-eater and, potentially, in other closely related species.

Keywords Cooperative breeding · Cross-species markers · European bee-eater · *Merops apiaster* · Microsatellite markers · Multiplex set

European bee-eaters (*Merops apiaster*) are a migratory bird species in the Meropidae family (order Coraciiformes). They breed in Southern Europe and the North of Africa and migrate to Southern Africa during winter (Fry 1984). Unlike many cooperatively breeding birds, European bee-eaters are a colonial species (Lessells et al. 1994). Colonies vary in size and can range from two up to several hundred breeding pairs (Fry 1984). Approximately 20 % of broods in this species obtain help from other individuals that have failed in their breeding attempts (Lessells 1990). Females have greater natal dispersal distances than males, although breeding-site fidelity is very common (Lessells et al. 1994). However, less is known about inter-colony dispersal (Dasmahapatra et al. 2004) and associated fine-scale genetic structure, features that may be important if we are to understand the evolution of various life-history traits such as cooperative breeding, coloniality and dispersal. Here, we characterise a set of polymorphic microsatellite loci, obtained via cross-species amplification, and combine these with existing European bee-eater loci (Dasmahapatra et al. 2004) and sexing markers (Dawson 2007) to create a powerful multiplex set. This multiplex set will facilitate the study of dispersal, parentage and kin selection in the European bee-eater and could potentially be of use in similar studies of closely related species.

A total of 42 microsatellite loci were tested for their utility in European bee-eaters, sampled on Susak Island, Croatia (45° 29' 26" N, 14° 19' 46" E). Eleven of these loci were previously developed for the European bee-eater (Dasmahapatra et al.

✉ Hannah L. Dugdale
h.dugdale@leeds.ac.uk

¹ Department of Animal and Plant Sciences, University of Sheffield, Western Bank Sheffield S10 2TN, UK

² Groningen Institute for Evolutionary Life Sciences, University of Groningen, PO Box 11103, 9700 CC Groningen, Netherlands

³ Department of Life Sciences, Imperial College London, Silwood Park Campus, Buckhurst Road, Ascot, Berks SL5 7PY, UK

⁴ Evolutionary Biology, Max Planck Institute for Ornithology, Eberhard Gwinner Strasse, 82319 Seewiesen, Germany

⁵ School of Biology, University of Leeds, Leeds LS2 9JT, UK

Table 1 Five microsatellite multiplex sets tested in adult European bee-eaters (*Merops apiaster*), including eight cross-species markers, seven markers isolated in European bee-eaters and two sex-typing markers

Multiplex	Marker	Source species & EMBL accession number & reference for primer set	Primer sequence (5'-3')	Fluoro label	N	N _a	Allele size range (bp)	H _O	H _E	HWE p value	Est. null allele freq.
1	Be2.31	<i>Merops apiaster</i> AJ630048 Dasmahapatra et al. 2004	F: CTTCAGGCAAGTGACCACAG	6-FAM	21	7	165–181	0.86	0.84	0.886	-0.021
			R: CAGAGGGACACCAGAGCTTC								
	Be2.33	<i>Merops apiaster</i> AJ630050 Dasmahapatra et al. 2004	F: CAGGAATGCTGTTGAACCTG	HEX	20	3	176–180	0.50	0.41	0.719	-0.135
			R: ACTGTGCCTTGCTCACATTG								
	Mor12	<i>Merops ornatus</i> DQ304696 Adcock et al. 2006	F: CATTCAAAAATACCCAGTGC	HEX	23	3	209–213	0.26	0.27	0.158	0.053
			R: AGACTGCTTCGTCTGTATGC								
	Mor20	<i>Merops ornatus</i> DQ304700 Adcock et al. 2006	F: CCACTTGTCTTTTGTCAAACC	6-FAM	21	6	206–216	0.67	0.74	0.283	0.042
			R: TCTCAGGTCAAAGCTGAAACC								
	Z002D	Dawson 2007	(Dawson 2007)	6-FAM	18 male 20 female	1 2	125, 125 (Z, Z) 114, 125 (W, Z)	0 -	0 -	-	-
			F: CTTTGTGAGTTTCCCTGTAGC								
Bb111-TG	<i>Bubo bubo</i> AF432096 (Bb111) Klein et al. 2009	R: ATCTAAGCAITAAAAATGCARAYCTT	HEX	18	6	188–200	0.56	0.63	0.380	0.065	
		F: AAGCAAGGACTTCTCCAG									
HvoB1-TTG	<i>Haliaeetus vociferoides</i> AJ620441 (HvoB1) Klein et al. 2009	R: TCTCAAATGGAAACAGAGAAAAGG	HEX	23	12	117–142	0.83	0.72	1.000	-0.112	
		F: GACAATGGCTATGAAAATAAATTAGGC									
TG04-061	<i>Taeniopygia guttata</i> CK235034 Dawson et al. 2010	R: AGAAGGGCAITGAAAGCACAC	HEX	18	8	210–218	0.67	0.75	0.087	0.055	
		(Dawson 2007)									
Z002D	Dawson 2007	(Dawson 2007)	6-FAM	18 male 20 female	1 2	125, 125 (Z, Z) 114, 125 (W, Z)	0 -	0 -	-	-	
		F: GATCACATTATCTGCATGIG									
Be3.24	<i>Merops apiaster</i> AJ630052 Dasmahapatra et al. 2004	R: TTATGAAAAGTCTACTTATATGTGTCC	HEX	23	10	141–163	0.78	0.87	0.243	0.043	
		F: GGATCTATAAACACATCTGCAT									
Be3.9	<i>Merops apiaster</i> AJ630053 Dasmahapatra et al. 2004	R: AAGGAATTACCTGCCCTTA	6-FAM	23	13	162–190	0.78	0.89	0.257	0.058	
		F: GGGTGTCTGTACACCTTGTAGC									
Mor10	<i>Merops ornatus</i> DQ304695 Adcock et al. 2006	R: AACGTGTCCCAAAATCTCC	HEX	23	3	198–202	0.61	0.54	0.220	-0.068	
		F: AACGTGTCCCAAAATCTCC									

Table 1 (continued)

Multiplex	Marker	Source species & EMBL accession number & reference for primer set	Primer sequence (5'-3')	Fluoro label	N	N _a	Allele size range (bp)	H _O	H _E	HWE p value	Est. null allele freq.
4	Z002D	Dawson 2007	(Dawson 2007)	6-FAM	18 male 20 female	1	125, 125 (Z, Z)	0	0	-	-
	Be1.29	<i>Merops apiaster</i> AJ630054 Dasmahapatra et al. 2004	F: TTTTCTCTGGGAGGTGGTTG R: GCTTGAAGGGGATTAIGATAGC	HEX	23	11	114, 125 (W, Z) 147-167	-	-	-	-
	Be19.2	<i>Merops apiaster</i> AJ630055 Dasmahapatra et al. 2004	F: GTCAAAGTGGGCTGTTGGAG R: AAGAGGGGCTACTTCCAAGC	NED	23	7	174-186	0.83	0.83	0.419	-0.013
	Be2.52	<i>Merops apiaster</i> AJ630049 Dasmahapatra et al. 2004	F: GGAGTCACTAGGCCCATCC R: TTTCGGAGGCAGTGTAAG	NED	18	13	214-240	0.83	0.90	0.431	0.024
5	Z002B	Dawson 2007	(Dawson 2007)	6-FAM	18 male 20 female	1	247, 247 (Z, Z) 241, 247 (W, Z)	0	0	-	-
	CAM-15	<i>Taeniopygia guttata</i> HG518773 Dawson et al. 2013	F: SGACGACTCCTTTAATTTCCC R: TTCTGACTTCCYCAGGTAACAC	6-FAM	21	7	261-273	0.81	0.82	0.519	-0.005
	TG13-017	<i>Taeniopygia guttata</i> CK313422 Dawson et al. 2010	F: GCTTTGCATCTTGCCCTAAA R: GGTAACACTACAACACTTCCAACCTCCT	HEX	22	4	209-217	0.50	0.58	0.712	0.072
	Z002B	Dawson 2007	(Dawson 2007)	6-FAM	18 male 20 female	1	247, 247 (Z, Z) 241, 247 (W, Z)	0	0	-	-

N is the number of genotyped adult European bee-eater (*Merops apiaster*) individuals from Susak Island, Croatia, N_a is the number of alleles, H_O is the observed heterozygosity, H_E is the expected heterozygosity and Est. null allele freq. is the estimated null allele frequency

2004), and we assessed these loci for their utility in our study population. The remaining 31 loci were selected based on their high utility in multiple other species (Dawson et al. 2010, 2013), their utility in species belonging to the same order as European bee-eaters ($N=7-23$; Appendix 1) or both. Five of these loci were originally isolated from the closely related Rainbow bee-eater (*Merops ornatus*; Adcock et al. 2006), and 26 were of high cross-species utility, including 24 that were previously characterised in a related but more distant Coraciiform species, the European roller (*Coracias garrulous*; Martín-Gálvez et al. 2014).

Blood samples (ca 50 μ l) were collected by puncturing the brachial vein of 38 European bee-eaters from 15 colonies on Susak Island in 2014. These samples were stored in 1 ml of 96 % ethanol in rubber-sealed screw-topped microfuge tubes. We extracted genomic DNA via an ammonium acetate precipitation method (Nicholls et al. 2000) and performed PCR amplification in a 2- μ l volume. Each PCR reaction contained approximately 15 ng of genomic DNA, 0.2 μ M of each primer (except for *Be2.52* [0.6 μ M], *Be3.9* [1.0 μ M], *CAM-15* [0.4 μ M], *Mor14* [0.6 μ M] and *TG13-017* [0.6 μ M]), HotStarTaq DNA polymerase, MgCl₂ and dNTPs supplied at the concentrations stated in the manufacturer's buffer (QIAGEN Multiplex PCR Master Mix). Forward primers were fluorescently labelled and PCR products were amplified using a DNA Engine Tetrad PTC-225 Peltier thermal cycler (MJ Research).

Loci of high cross-species utility and those tested in the European roller were amplified in singleplexes across all 38 individuals with the following PCR profile: 95 °C for 15 min, followed by 35 cycles of 94 °C for 30 s, 56 °C for 90 s and 72 °C for 60 s and a final step of 60 °C for 30 min. The Rainbow and European bee-eater loci were initially assessed in multiplex reactions (designed using Multiplex Manager 1.2) in seven adults that were not close relatives ($R \leq 0.126$; GenAlEx 6.41, Peakall and Smouse 2006) with the same PCR constituents, including 0.2 μ M of each primer (except for *Be2.52* [0.6 μ M], *Be3.9* [1.0 μ M] and *Mor14* [0.6 μ M]), and the same PCR profile. PCR products were separated on an ABI 3730 DNA Analyser with an ABI ROX500 size standard and alleles scored using GENEMAPPER 3.7 (Applied Biosystems).

We found 20 loci to be polymorphic in seven to 38 individuals. Five loci were monomorphic and 17 failed to amplify a product (Appendix 1). The five monomorphic loci in this population may be variable and so of utility in other populations. Locus *Be72* is the same as *Be3.9*, which we tested under the latter name and found polymorphic (Adcock et al. 2006). Polymorphic loci were amplified as multiplex PCRs across all 38 individuals using the PCR profile stated above. Each multiplex set included a marker of known utility for sexing European bee-eaters; *Z002B* or *Z002D* (Dawson 2007). Loci were then characterised across a sub-set of 23 European bee-eaters, by selecting one male and one female adult from each colony to minimise overlapping generation and social structure effects. All

20 loci were autosomal as heterozygotes occurred in both sexes (18 males and 20 females assessed). One autosomal locus (*TG01-000*) amplified successfully in singleplex but failed to amplify in multiplex PCR (Appendix 1). Another locus (*Be48*) was polymorphic with only three alleles and little genetic variation in the Susak Island individuals ($H_O=0.09$, Appendix 1). Despite the relatively low variability among Susak Island individuals, this locus may be suitable for population structure analyses as heterozygosity was higher ($H_O=0.29$) in a French European bee-eater population (Dasmahapatra et al. 2004).

We used data from the 23 adult European bee-eaters to test each locus for deviation from Hardy-Weinberg Equilibrium (HWE) and for linkage disequilibrium among groups of loci (GENEPOP 4.2; Rousset 2008). Three loci (*Be2.16*, *Mor14* and *Mor15*) deviated from HWE and were not included in the final multiplex set (Appendix 1). Loci *Be2.16* and *Mor15* also displayed high null allele frequency estimates (>0.10 ; CERVUS 3.0.03, Kalinowski et al. 2007). After correcting for multiple testing (Benjamini and Hochberg 1995), three pairs of loci (*TG04-061* & *CAM-15*, *Mor10* & *Be19.2* and *Mor10* & *Bb111-TG*) displayed linkage disequilibrium ($p < 0.05$). We therefore omitted data from *CAM-15* and *Mor10* from the analyses of relatedness. The final 15 autosomal loci (eight cross-species loci combined with seven loci isolated in European bee-eaters) were checked and confirmed unique using BLAST software (Altschul et al. 1997). These 15 loci, optimised in five multiplexes, will be useful in future studies, especially when combined with the two sex-typing markers (Table 1). In particular, the marker set will facilitate both fine-scale and large-scale population genetic structure analyses, and facilitate parentage assignment and relatedness estimation for kin selection studies. More specifically, this multiplex set will provide an efficient and economical method for investigating the social dynamics of European bee-eaters and, potentially, closely related species.

Acknowledgments Molecular work was carried out in the Molecular Ecology Laboratory at the University of Sheffield, UK. Simon Weigl and Liviu Parau assisted with sample collection. This work was carried out under licenses from the Croatian Ministry of Environmental and Nature Protection and the Croatian Institute for Ornithology.

Compliance with ethical standards

Funding GB was funded by a Genes and Development Summer Studentship (Genetics Society). DAD was funded by the Natural Environment Research Council (NERC), UK. HLD and NdR were funded by a NERC fellowship (NE/I021748/1). JS was funded by the Volkswagen Foundation. SAK was funded by a VENI-fellowship awarded by the Netherlands Organisation for Scientific Research (NWO 863.13.017). Fieldwork was funded by the Nicolaas Mulerius Foundation (University of Groningen) and by the Ecology Fund (Royal Netherlands Academy of Arts and Sciences; KNAW UPS/297/Eco/1414J).

Ethical approval All applicable international, national and/or institutional guidelines for the care and use of animals were followed.

This article does not contain any studies with human participants performed by any of the authors.

Appendix

Appendix 1 Details of the 42 microsatellite markers tested in the European bee-eater (*Merops apiaster*). Markers in italics and underlined were included in the final multiplex set

Marker	Source species & EMBL accession number & locus name (when different)	Reference for primer set	Reason for testing	Fluoro label	Plex used in initial test	N	N_a	Allele size range (bp)	Outcome
<u>ApCo46-ZEST</u>	<i>Aphelocoma coerulescens</i>	Stenzler et al. 2002	1, 2	6-FAM	Singleplex	7	–	–	PCR failed
ApCo80-CEST	AF520885 (ApCo46) <i>Aphelocoma coerulescens</i>	Stenzler et al. 2002	1, 2	HEX	Singleplex	7	–	–	PCR failed
<u>Bb111-TG</u>	AF520879 (ApCo80) <i>Bubo bubo</i> (AF432096, Bb111), <i>Taeniopygia guttata</i> , <i>Gallus gallus</i> <i>Merops apiaster</i>	Klein et al. 2009	1, 2	HEX	Singleplex	18	6	188–200	Polymorphic, see Table 1
<u>Be1.29</u>	AJ630054 <i>Merops apiaster</i>	Dasmahapatra et al. 2004	3	HEX	Multiplex	23	11	147–167	Polymorphic, see Table 1
<u>Be19.2</u>	<i>Merops apiaster</i>	Dasmahapatra et al. 2004	3	NED	Multiplex	23	7	174–186	Polymorphic, see Table 1
Be2.16	AJ630055 <i>Merops apiaster</i>	Dasmahapatra et al. 2004	3	NED	Multiplex	23	5	189–195	Polymorphic but rejected due to high estimated null allele frequency (0.251) and deviation from HW (p = 0)
<u>Be2.31</u>	AJ630047 <i>Merops apiaster</i>	Dasmahapatra et al. 2004	3	6-FAM	Multiplex	21	7	165–181	Polymorphic, see Table 1
<u>Be2.33</u>	AJ630048 <i>Merops apiaster</i>	Dasmahapatra et al. 2004	3	HEX	Multiplex	20	3	176–180	Polymorphic, see Table 1
Be2.46	AJ630050 <i>Merops apiaster</i>	Dasmahapatra et al. 2004	3	6-FAM	Multiplex	7	1	187	Monomorphic
<u>Be2.52</u>	AJ630051 <i>Merops apiaster</i>	Dasmahapatra et al. 2004	3	NED	Multiplex	18	13	214–240	Polymorphic, see Table 1
Be24	AJ630049 <i>Merops apiaster</i>	Dasmahapatra et al. 2004	3	HEX	Multiplex	8	–	–	PCR failed; did not amplify as singleplex or multiplex
<u>Be3.24</u>	AJ630044 <i>Merops apiaster</i>	Dasmahapatra et al. 2004	3	HEX	Multiplex	23	10	141–163	Polymorphic, see Table 1

Appendix 1 (continued)

Marker	Source species & EMBL accession number & locus name (when different)	Reference for primer set	Reason for testing	Fluoro label	Plex used in test	N	N_a	Allele size range (bp)	Outcome
		Dasmahapatra et al. 2004							
<u>Be3.9</u>	AJ630052 <i>Merops apiaster</i>	Dasmahapatra et al. 2004	3	6-FAM	Multiplex	23	13	162–190	Polymorphic, see Table 1
Be48	AJ630053 <i>Merops apiaster</i>	Dasmahapatra et al. 2004	3	6-FAM	Multiplex	23	2	151–153	Polymorphic but rejected due to low heterozygosity ($H_o = 0.09$)
CAM-06	AJ630045 <i>Taeniopygia guttata</i> (HG518764), <i>Gallus gallus</i>	Dawson et al. 2013	1	HEX	Singleplex	7	–	–	PCR failed
CAM-12	<i>Taeniopygia guttata</i> (HG518770), <i>Gallus gallus</i>	Dawson et al. 2013	1, 2	6-FAM	Singleplex	7	–	–	PCR failed
CAM-13	<i>Taeniopygia guttata</i> (HG518771), <i>Gallus gallus</i>	Dawson et al. 2013	1, 2	HEX	Singleplex	7	–	–	PCR failed
<u>CAM-15</u>	<i>Taeniopygia guttata</i> (HG518773), <i>Gallus gallus</i>	Dawson et al. 2013	1, 2	6-FAM	Singleplex	21	7	261–273	Polymorphic, see Table 1
CAM-16	<i>Taeniopygia guttata</i> (HG518774), <i>Gallus gallus</i>	Dawson et al. 2013	1, 2	HEX	Singleplex	14	–	–	PCR failed
CAM-18	<i>Taeniopygia guttata</i> (HG518776), <i>Gallus gallus</i>	Dawson et al. 2013	1, 2	HEX	Singleplex	7	–	–	PCR failed
CAM-20	<i>Taeniopygia guttata</i> (HG518778), <i>Gallus gallus</i>	Dawson et al. 2013	2	HEX	Singleplex	7	–	–	PCR failed
CAM-24	<i>Taeniopygia guttata</i> (HG518782), <i>Gallus gallus</i>	Dawson et al. 2013	1, 2	HEX	Singleplex	7	–	–	PCR failed
<u>HvoB1-TTG</u>	<i>Haliaeetus vociferoides</i> (AJ620441, HvoB1), <i>Taeniopygia guttata</i> , <i>Gallus gallus</i>	Klein et al. 2009	1, 2	HEX	Singleplex	23	12	117–142	Polymorphic, see Table 1
<u>Mor10</u>	<i>Merops ornatus</i> DQ304695	Adcock et al. 2006	2	HEX	Multiplex	23	3	198–202	Polymorphic, see Table 1

Appendix 1 (continued)

Marker	Source species & EMBL accession number & locus name (when different)	Reference for primer set	Reason for testing	Fluoro label	Plex used in initial test	N	N_a	Allele size range (bp)	Outcome
<u>Mor12</u>	<i>Merops ornatus</i> DQ304696	Adcock et al. 2006	2	HEX	Multiplex	23	3	209–213	Polymorphic, see Table 1
Mor14	<i>Merops ornatus</i>	Adcock et al. 2006	2	NED	Multiplex	23	5	195–207	Polymorphic but rejected due to deviation from HWE ($p = 0.044$)
Mor15	DQ304697 <i>Merops ornatus</i>	Adcock et al. 2006	2	NED	Multiplex	22	9	158–180	Polymorphic but rejected due to high estimated null allele frequency (0.254) and deviation from HWE ($p = 0.002$)
<u>Mor20</u>	DQ304698 <i>Merops ornatus</i> DQ304701	Adcock et al. 2006	2	6-FAM	Multiplex	21	6	206–216	Polymorphic, see Table 1
Pte24-CEST	<i>Pomatostomus temporalis</i> DQ234870 (Pte24)	Blackmore et al. 2006	2	HEX	Singleplex	14	–	–	PCR failed
SAP47-ZEST	<i>Sayornis phoebe</i> AY823673 (SAP47)	Beheler et al. 2007	1, 2	6-FAM	Singleplex	7	–	–	PCR failed
Tc.11B4E-CEST	<i>Telespiza cantans</i> AF036266 (Tc.11B4E)	Tarr et al. 1998	1, 2	6-FAM	Singleplex	7	1	153	Monomorphic
TG01-000	<i>Taeniopygia guttata</i> (CK314156), <i>Gallus gallus</i>	Dawson et al. 2010	1, 2	6-FAM	Singleplex	14	2	205–207	Polymorphic, but, although it amplified in singleplex, failed in multiplex
TG01-040	<i>Taeniopygia guttata</i> (DV576233), <i>Gallus gallus</i>	Dawson et al. 2010	1, 2	6-FAM	Singleplex	7	1	288	Monomorphic
TG02-078	<i>Taeniopygia guttata</i> (CK305233), <i>Gallus gallus</i>	Dawson et al. 2010	1, 2	HEX	Singleplex	14	–	–	PCR failed
TG03-002	<i>Taeniopygia guttata</i> (DV575298), <i>Gallus gallus</i>	Dawson et al. 2010	1, 2	6-FAM	Singleplex	14	1	119	Monomorphic
TG03-098	<i>Taeniopygia guttata</i> (DV573670), <i>Gallus gallus</i>	Dawson et al. 2010	1, 2	HEX	Singleplex	14	–	–	PCR failed

Appendix 1 (continued)

Marker	Source species & EMBL accession number & locus name (when different)	Reference for primer set	Reason for testing	Fluoro label	Plex used in initial test	N	N_a	Allele size range (bp)	Outcome
TG04-012	<i>Taeniopygia guttata</i> (CK306810), <i>Gallus gallus</i>	Dawson et al. 2010	1, 2	HEX	Singleplex	–	–	–	PCR failed
<u>TG04-061</u>	<i>Taeniopygia guttata</i> (CK235034), <i>Gallus gallus</i>	Dawson et al. 2010	1, 2	HEX	Singleplex	18	8	210–218	Polymorphic, see Table 1
TG05-053	<i>Taeniopygia guttata</i> (CK314425), <i>Gallus gallus</i>	Dawson et al. 2010	1, 2	6-FAM	Singleplex	14	–	–	PCR failed
TG11-011	<i>Taeniopygia guttata</i> (CK308096), <i>Gallus gallus</i>	Dawson et al. 2010	1, 2	6-FAM	Singleplex	7	–	–	PCR failed
<u>TG13-017</u>	<i>Taeniopygia guttata</i> (CK313422), <i>Gallus gallus</i>	Dawson et al. 2010	1, 2	HEX	Singleplex	22	4	209–217	Polymorphic, see Table 1
Tgu06	<i>Taeniopygia guttata</i> CK307697	Slate et al. 2007	1, 2	HEX	Singleplex	7	1	172	Monomorphic

N is the number of European bee-eater (*Merops apiaster*) individuals tested/genotyped from Susak Island, Croatia, N_a is the number of alleles. Please note, many of these markers were donated from earlier studies and failure to amplify could be a result of the age of the primers.

Reason for testing:

- 1: Marker of known high cross-species utility (Dawson et al. 2010, 2013, DAD unpublished data)
- 2: Marker possessed proven utility in a related Coraciiform species: Rainbow bee-eater (*Merops ornatus*; Adcock et al. 2006) or European roller (*Coracias garrulus*; Martín-Gálvez et al. 2014)
- 3: Locus isolated from study species European bee-eater (*Merops apiaster*, Dasmahapatra et al. 2004)

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Adcock GJ, Hodges K, Boland CRJ, Cockburn A, Ebert D, Heinsohn R (2006) Microsatellite loci for behavioural studies of rainbow bee-eaters (*Merops ornatus*: Aves). *Mol Ecol Notes* 6:734–736
- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and psi-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25:3389–3402
- Beheler AS, Fike JA, Rhodes OE Jr (2007) Eight new polymorphic microsatellite loci from the eastern phoebe (*Sayornis phoebe*). *Conserv Genet* 8:1259–1261
- Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Ser B* 57:289–300
- Blackmore CJ, Adcock GJ, Ebert D, Heinsohn R (2006) Microsatellite loci for population and behavioural studies of grey-crowned babblers (*Pomatostomus temporalis*: Aves). *Mol Ecol Notes* 6:412–414
- Dasmahapatra KK, Lessells CM, Mateman AC, Amos W (2004) Microsatellite loci in the European bee-eater, *Merops apiaster*. *Mol Ecol Notes* 4:500–502
- Dawson DA (2007) Genomic analysis of passerine birds using conserved microsatellite loci. PhD Thesis, University of Sheffield, UK
- Dawson DA, Horsburgh GJ, Küpper C, Stewart IRK, Ball AD, Durrant KL, Hansson B, Bacon I, Bird S, Klein Á, Lee J-W, Martín-Gálvez D, Simeoni M, Smith G, Spurgin LG, Burke T (2010) New methods to identify conserved microsatellite loci and develop primer sets of high utility—as demonstrated for birds. *Mol Ecol Resour* 10:475–494
- Dawson DA, Ball AD, Spurgin LG, Martín-Gálvez D, Stewart IR, Horsburgh GJ, Potter J, Molina-Morales M et al (2013) High-utility conserved avian microsatellite markers enable parentage and population studies across a wide range of species. *BMC Genomics* 14:176
- Fry CH (1984) The bee-eaters. Buteo books. T & A D Poyser, UK
- Kalinowski ST, Taper ML, Marshall TC (2007) Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Mol Ecol* 16:1099–1106
- Klein A, Horsburgh GJ, Küpper C, Major A, Lee PLM, Hoffmann G, Mátics R, Dawson DA (2009) Microsatellite markers characterized in the barn owl (*Tyto alba*) and of high utility in other owls (*Strigiformes*: AVES). *Mol Ecol Resour* 9:1512–1519
- Lessells CM (1990) Helping at the nest in European bee-eaters: who helps and why? In: Population biology of passerine birds, an integrated approach, NATO ASI series. Springer-Verlag, Berlin, pp 357–368
- Lessells CM, Avery MI, Krebs JR (1994) Nonrandom dispersal of kin: why do European bee-eater (*Merops apiaster*) brothers nest close together? *Behav Ecol* 5:105–113
- Martín-Gálvez D, Molina-Morales M, Dawson DA, Parejo D, Martínez JG, Avilés JM (2014) Characterization of 28 microsatellite loci in the European roller *Coracias garrulus* (*Coraciidae*, AVES). *Eur J Wildl Res* 60:843–850
- Nicholls JA, Double MC, Rowell DM, Magrath RD (2000) The evolution of cooperative and pair breeding in thornbills *Acanthiza (Pardalotidae)*. *J Avian Biol* 31:165–176
- Peakall R, Smouse PE (2006) GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Mol Ecol Notes* 6:288–295
- Rousset F (2008) GENEPOP' 007: a complete re-implementation of the GENEPOP software for Windows and Linux. *Mol Ecol Resour* 8:103–106
- Slate J, Hale MC, Birkhead TR (2007) Simple sequence repeats in zebra finch (*Taeniopygia guttata*) expressed sequence tags: A new resource for evolutionary genetic studies of passerines. *BMC Genomics* 8:52
- Stenzler LM, Fitzpatrick JW (2002) Isolation of microsatellite loci in the Florida Scrub-Jay *Aphelocoma coerulescens*. *Mol Ecol Notes* 2:547–550
- Tarr CL, Conant S, Fleischer RC (1998) Founder events and variation at microsatellite loci in an insular passerine bird, the Laysan finch (*Telespiza cantans*). *Mol Ecol* 7:719–731